Printed Pages:02 Sub Code:KOE-067
Paper Id: 236488 Roll No.

B.TECH. (SEM VI) THEORY EXAMINATION 2022-23 BASICS OF DATA BASE MANAGEMENT SYSTEM

Time: 3 Hours Total Marks: 100

Note: Attempt all Sections. If require any missing data; then choose suitably.

SECTION A

1. Attempt *all* questions in brief.

 $2 \times 10 = 20$

- a. Define Database, DBMS, and Metadata.
- b. Discuss the problems with File systems.
- Define Instance and Domain.
- d. Define Recursive Relationship with example.
- e. Define 3NF and BCNF.
- f. Define the term Data Dictionary.
- g. List and describeSQL Logical Operators in brief.
- h. Describe DROP command with options.
- i. Define Two Phase Commit protocol.
- j. Define a checkpoint.

SECTION B

2. Attempt any *three* of the following:

10x3 = 30

- a. Define Data Abstraction and discuss levels of Abstraction.
- b. Explain about different types of integrity constraints with examples.
- c. Illustrate Group by and Having clauses with examples.
- d. Define decomposition and how does it address redundancy? Discuss the problems that may be caused by the use of decompositions?
- e. Discuss Serializability in detail.

SECTION C

3. Attempt any *one* part of the following:

10x1=10

- a. Describe the Structure of DBMS. Write about views and updates on views.
- b. Distinguish strong entity set with weak entity set. Draw an ER diagram to illustrate weak entity set?

4. Attempt any *one* part of the following:

10x1=10

- Define Join. Explain different types of joins. a.
- b. Discuss different types of aggregate operators with examples in SQL?

5. Attempt any *one* part of the following:

10x1=10

- a. Define functional dependencies and relation of primary keys with FD's?
- b. Illustrate Multivalued dependencies and Fourth normal form with example.

6. Attempt any *one* part of the following:

10x1=10

- Explain ACID properties and Illustrate them through examples. a.
- b. Describe Validation-based locking protocols. Discuss in detail Multiple Granularity.
- 7. Attempt any *one* part of the following:

10x1=10

- a.
- Explain Armstrong's inference rules?
 Suppose we are given relation R with attributes A, B, C, D, E, F, and the FDs, A→BC, B→E, CD→EF. Prove that FD: AD→F also holds in R. b.